

Stasis Points and Approximating Two-Cycles

Stewart D. Johnson
Department of Mathematics and Statistics
Williams College

February 1, 2008

Abstract

Generically, every fixed point for the differential inclusion $x' \in \text{Convex Hull}\{f_1(x), f_2(x)\}$ can be approximated by an arbitrarily small two-cycle for the inclusion $x' \in \{f_1(x), f_2(x)\}$, where f_1, f_2 are a C^1 flows on $\mathbb{R}^n$.

1 Two-Flows, Stasis Points and Two-Cycles

Consider a controlled system that switches freely between two autonomous flows $x' = f_1(x)$ and $x' = f_2(x)$ where $x \in \mathbb{R}^n$ and $f_i : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is C^1 . This is the differential inclusion

$$x' \in \{f_1(x), f_2(x)\}.$$

Trajectories are piecewise C^1 functions $x(t)$ satisfying this inclusion.

A **stasis point** is a point x at where the flows are anti-parallel:

$$f_1(x) \cdot f_2(x) = -|f_1(x)| \cdot |f_2(x)|.$$

The set of stasis points is the set of possible fixed points for the relaxed differential inclusion

$$x' \in \text{Convex Hull}(\{f_1(x), f_2(x)\}).$$

A **two-cycle** $x(t)$ is a periodic trajectory $x(t) = x(\delta + t)$ with $0 < \kappa < \delta$ and

$$\begin{aligned} x' &= f_1(x) & \text{for } 0 \leq t < \kappa \\ \text{and } x' &= f_2(x) & \text{for } \kappa \leq t < \delta \end{aligned}$$

It will be shown that neighborhoods of stasis points generically contain two-cycles.

2 Loops in the Action

An C^2 **action** of $\mathbb{R}$ on $\mathbb{R}^n$ is a map $\Phi : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$ which is C^2 in all variables. A **loop** is a point and time x_0, t_0 with $\Phi(x_0, t_0) = \Phi(x_0, t_0 + \delta)$ for some $\delta > 0$.

A point x_0 is **stationary** at time t_0 if $\frac{\partial}{\partial t}\Phi(x_0, t_0) = \vec{0}$. Note that a point that remains stationary for some open time interval, $\frac{\partial}{\partial t}\Phi(x_0, t) \equiv \vec{0}$ for $\alpha < t < \beta$, is a (degenerate) loop of all sufficiently small periods.

The following theorem states that actions with stationary points generically have small loops nearby.

Theorem 1 *Let $\Phi : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}^n$ be a C^2 action. Suppose $\exists x_0, t_0$ with $\frac{\partial}{\partial t}\Phi(x_0, t_0) = \vec{0}$. If*

$$\frac{\partial^2}{\partial x \partial t}\Phi(x_0, t_0)$$

is non-singular then $\forall \epsilon, \exists z_0, \tau_0, \delta > 0$ with $\Phi(z_0, \tau_0) = \Phi(z_0, \tau_0 + \delta)$ and

$$|\Phi(z_0, t) - \Phi(x_0, t_0)| < \epsilon$$

for $\tau_0 \leq t < \tau_0 + \delta$.

Proof:

Without loss of generality we take $x_0 = \vec{0}$ and $t_0 = 0$.

Let

$$G(x, \delta) = \begin{cases} \frac{1}{2\delta} (\Phi(x, \delta) - \Phi(x, -\delta)) & \text{if } \delta \neq 0 \\ \frac{\partial}{\partial t} \Phi(x, 0) & \text{if } \delta = 0 \end{cases}$$

So G is C^1 at $x = \vec{0}, \delta = 0$, and $\frac{\partial G}{\partial x}(\vec{0}, 0) = \frac{\partial^2 \Phi}{\partial x \partial t}(\vec{0}, 0)$ is non-singular by assumption. Therefore the Implicit Function Theorem applies, and since $G(\vec{0}, 0) = \vec{0}$ there exist solutions x to $G(x, \delta) = \vec{0}$ for δ sufficiently small.

For $\delta \neq 0$, a point x with $G(x, \delta) = \vec{0}$ has $\Phi(x_0, \delta) = \Phi(x_0, -\delta)$, which is a loop. ■

3 Two-Cycle Approximation to Stasis

It is known that relaxed solutions to differential inclusions can be arbitrarily approximated by non-relaxed solutions [1, 2]. Here it is shown that in the case of autonomous two-flows and relaxed fixed points, the approximating trajectory can be taken as a two-cycle. For flows f_1, f_2 , the idea is to take $\mathbb{R}^n$ modulo the trajectories of f_1 , and consider the remaining (non-autonomous) action of f_2 on $\mathbb{R}^{n-1}$. Loops in this resulting action will pull back to two-cycles in the original system.

Theorem 2 Consider two C^1 flows $f_1, f_2 : \mathbb{R}^n \rightarrow \mathbb{R}^n$, and a stasis point x_0 with $k_1 f_1(x_0) + k_2 f_2(x_0) = \vec{0}$, $k_i > 0$. If $\frac{d}{dx} (k_1 f_1 + k_2 f_2)$ is non-singular at x_0 , then there exist two-cycles near x_0 .

Proof: Without loss of generality we take $x_0 = \vec{0}$. Since f_1 is C^1 and non-zero at near $\vec{0}$, there is a local diffeomorphism $\rho : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with $\frac{d\rho}{dx} \cdot f_1 \equiv (1, 0, \dots, 0)$ near $\vec{0}$. Then $g_2 = \frac{d\rho}{dx} \cdot f_2 \circ \rho^{-1}$ is a C^1 flow with $g_2(\vec{0}) = (-\frac{k_1}{k_2}, 0, \dots, 0)$.

We construct the $\mathbb{R}^{n-1}$ action Φ as follows: given $y = (a_1, \dots, a_{n-1}) \in \mathbb{R}^{n-1}$ let $x(t) = (x_1(t), \dots, x_n(t))$ be the trajectory under the flow g_2 with $x(0) =$

$(0, a_1, \dots, a_{n-1}) \in \mathbb{R}^n$. Taking $\Phi(y, t) = (x_2(t), \dots, x_n(t))$ defines a C^2 action on $\mathbb{R}^{n-1}$ with $\frac{\partial}{\partial t}\Phi(\vec{0}, 0) = \vec{0}$. It is straightforward to check that non-singularity of $\frac{d}{dx}(\alpha f_1 + \beta f_2)$ at $\vec{0} \in \mathbb{R}^n$ implies that of $\frac{\partial^2}{\partial x \partial t}\Phi$ at $(\vec{0}, 0) \in \mathbb{R}^{n-1} \times \mathbb{R}$.

By Theorem 1, for all $\epsilon > 0$ there exist $z = (z_1, \dots, z_{n-1})$ near $\vec{0} \in \mathbb{R}^{n-1}$ with $\Phi(z, \alpha) = \Phi(z, \beta)$ for some $|\alpha|, |\beta| < \epsilon$, $\alpha \neq \beta$. This corresponds to a trajectory $x(t)$ in under g_2 on $\mathbb{R}^n$ with $x(0) = (0, z_1, \dots, z_{n-1})$. Now $x(\alpha)$ and $x(\beta)$ are equal except for their first coordinate, which cannot be zero for sufficiently small ϵ because g_2 is non-zero near $\vec{0}$. Thus $x(\alpha) - x(\beta)$ is parallel to the constant flow g_1 and connecting these points with a segment of the g_1 flow completes the two-cycle. This maps to a two-cycle for f_1 and f_2 under ρ^{-1} . ■

4 Three Questions

For $\mathbb{R}^2$, every two cycle contains a stasis point [3]. Is a similar statement true for $\mathbb{R}^n$?

Given a pair of flows f_1, f_2 , the stasis points generically form one dimensional curves. For $n = 2$, an approximating two cycle will cut across this curve twice, and so the set of approximating two-cycles is parameterized by pairs of nearby points on the stasis curve [3]. Is there a similar organization of two-cycles in $\mathbb{R}^n$?

Finally, a natural question is to extend this to N -flows. That is, for a differential inclusion $x' \in \{f_1(x), f_2(x), f_3(x)\}$, can a relaxed fixed point always be approximated by a three-cycle?

References

- [1] A. F. Filippov, Differential equations with discontinuous righthand sides, Kluwer Academic, Dordrecht, The Netherlands, 1988.
- [2] B. Ingalls, E.D. Sontag, Y. Wang, *An infinite time relaxation theorem for differential inclusions*, Proc. of AMS, Vol 131, **2**, Pages 487-499, 2002.
- [3] S.D. Johnson, *Stasis and two-cycles*, SIAM J of Op Control, to appear.